

## Production of quality additive cereal forage crops on livestock digestibility by the impact of irrigation using distillery spentwash

Bhuvan P, Chandraju S

Department of Studies in Sugar Technology, Sir M. Vishweshwaraya Postgraduate Centre, University of Mysore, Tubinakere, Mandya, Karnataka, India.

### Abstract

Cultivation of cereal forages namely; Maize (*Zea mays*), Bajra (*Pennisetum Glaucum*) and Oats (*Avena sativa*) was made by irrigation with distillery spentwash of different proportions. The distillery spentwash i.e., primary treated spentwash (PTSW), 1:1, 1:2 and 1:3 distillery spentwash were analysed for plant nutrients such as nitrogen, phosphorous, potassium (NPK) and other physical and chemical parameters. The plants were cultivated by irrigation with raw water (RW), 1:1, 1:2 and 1:3 distillery spentwash in the prepared pots. The impact of distillery spentwash on proximate principles for quality forage (Crude protein, Neutral detergent fibre, Acid detergent fibre and Total digestible nutrient) i.e., forage digestibility for livestock were analysed. It was observed that good nutrients uptake in case of 1:3 spentwash and requirements of livestock digestibility components were observed when compared with 1:1, 1:2 spentwash and raw water irrigations. This could be due to the maximum absorption of NPK by plants at more diluted condition of spentwash. This concludes that the diluted spentwash can be conveniently used for the effective cultivation without using any external fertilizers. Hence, spentwash serves as a liquid fertilizer, eco-friendly irrigation medium and without adverse effect on environment and soil.

**Keywords:** Distillery spentwash, Maize, Bajra, Oats, Irrigation, Proximate principle

### 1. Introduction

Molasses is one of the important by-products of sugar industry which is the chief source for the production of alcohol in distilleries by fermentation method. Nearly 10-12 litres of spentwash are discharged for every litre of rectified spirit produced and is known as raw spent wash (RSW), which is characterized by high biological oxygen demand (BOD: 5000-8000mg/l) and chemical oxygen demand (COD: 25000-30000mg/l) (Joshi, 1994) [12], undesirable colour and foul odour. The discharge of spentwash into open field or water bodies result in environmental, soil & water pollution. Hence discharge of spent wash is a great problem. The RSW is highly acidic and consists of easily oxidisable organic matter with very high BOD and COD (Patil, 1987) [17]. The spentwash is rich in organic carbon & plant nutrients (Ramadurai and Gearard, 1994) [20]. Since it is from plant source extract it contains negligible heavy metals & other toxic substances (Eyini *et al.*, 1990) [27]. Meanwhile it is rich in plant essential nutrients it can be used in agriculture so the problem of disposal becomes easy along with the utilisation of nutrients by plants. It also helps to utilise spentwash in a proper method to avoid adverse effects on the environment. Its application to soil has been reported to be beneficial to increase sugar cane (Zalwadia, 1997) [26], rice (Devarajan and Oblisami, 1995) [10], wheat and rice yield (Pathak *et al.*, 1998) [18], quality of groundnut (Amar Singh *et al.*, 2003) [11] and physiological response of soybean (Ramana *et al.*, 2000) [21].

Diluted spent wash could be used for irrigation purpose without adversely affecting soil fertility (Kaushik *et al.*, 2005; Kuntal *et al.*, 2004; Raverkar *et al.*, 2000) [13, 14], seed germination and crop productivity (Ramana *et al.*, 2001). The diluted spent wash irrigation improved the physical and chemical properties of the

soil and further increased soil micro flora (Devarajan, 1994; Kaushik *et al.*, 2005; Kuntal *et al.*, 2004) [11, 13, 14]. Twelve pre sowing irrigations with the diluted spent wash had no adverse effect on the germination of maize but improved the growth and yield (Singh and Raj Bahadur, 1998) [22]. Diluted spent wash increases the growth of shoot length, leaf number per plant, leaf area and chlorophyll content of peas (Ravi and Srivastava, 1990). Increased concentration of spent wash causes decreased seed germination, seedling growth and chlorophyll content in sunflowers (*Helianthus annuus*) and the spent wash could be safely used for irrigation purpose at lower concentration (Rajendra, 1990; Ramana *et al.*, 2001) [23].

The spent wash contained an excess of various forms of cations and anions, which are injurious to plant growth and these constituents should be reduced to beneficial level by diluting the spent wash, which can be used as a substitute for chemical fertilizer (Sahai *et al.*, 1983) [24]. The spent wash could be used as a complement to mineral fertilizer to sugarcane (Chares, 1985) [4]. The spent wash contained N, P, K, Ca, Mg and S and thus valued as a fertilizer when applied to soil through irrigation with water (Samual, 1986) [25]. The application of diluted spent wash increased the uptake of Zinc(Zn), Copper(Cu), Iron(Fe) and Manganese(Mn) in maize and wheat as compared to control and the highest total uptake of these were found at lower dilution levels than at higher dilution levels (Pujar,1995) [19]. Mineralization of organic material as well as nutrients present in the spent wash was responsible for increased availability of plant nutrients. Diluted spent wash increases the uptake of nutrients, height, growth and yield of leafy vegetables (Chandraju *et al.*, 2007; Basvaraju and Chandraju, 2008) [2, 3, 5, 6, 7, 8, 9], nutrients of cabbage and mint leaf (Chandraju *et al.*, 2008) [2, 3, 6, 7, 8, 9], nutrients of top vegetable (Basvaraju and Chandraju, 2008) [2, 3,

6, 7, 8, 9], pulses, condiments and root vegetables (Chandraju *et al.*, 2008) [2, 3, 6, 7, 8, 9]. However, not much information is available on the impact of distillery spent wash on the production of quality additive cereal forage crops. Therefore, the present investigation was carried out to investigate the impact of irrigation of different concentration of spentwash on the production of quality additive cereal forage crops on livestock digestibility.

**2. Materials and Methods**

Physio-chemical parameters and amount of nitrogen (N), potassium (K), phosphorous (P) and sulphur (S) present in the primary treated diluted spentwash (1:1, 1:2, 1:3 SW) were analysed by standard methods. The PTSW was used for irrigation with a dilution of 1:1, 1:2 & 1:3. A composite soil sample collected prior to spentwash irrigation was air-dried, powdered and analysed for physio-chemical properties. The forage plants selected for the present investigation were Maize, Bajra and Oats seeds which were sowed in different pots [25.5cm (h), 45.5cm (dia)] and irrigated by applying 0.75 to 1 lit/pot (depending upon the climatic condition) with raw water (RW), 1:1 SW, 1:2 SW and 1:3 SW at the dosage of once a week and rest of the period with raw water as required. At the maturity time, forage samples were harvested air dried and proximate

principles on forage quality were analysed.

**Table 1:** Characteristics of experimental soil

Parameters	Values
Coarse sand <sup>c</sup>	8.99
Fine sand <sup>c</sup>	41.06
Slit <sup>c</sup>	25.87
Clay <sup>c</sup>	21.80
pH (1:2 soln)	8.32
Electrical conductivity <sup>a</sup>	562
Organic carbon <sup>c</sup>	0.98
Available Nitrogen <sup>b</sup>	392
Available Phosphorous <sup>b</sup>	239
Available Potassium <sup>b</sup>	99
Exchangeable Calcium <sup>b</sup>	163
Exchangeable Magnesium <sup>b</sup>	251
Exchangeable Sodium <sup>b</sup>	119
Available Sulphur <sup>b</sup>	296
DTPA Iron <sup>b</sup>	201
DTPA Manganese <sup>b</sup>	210
DTPA Copper <sup>b</sup>	9
DTPA Zinc <sup>b</sup>	62

Units: a-µS, b- mg/L, c-%

**Table 2:** Chemical characteristics of distillery Spentwash

Chemical parameters	PTSW	1:1 PTSW	1:2 PTSW	1:3 PTSW
PH	7.52	7.60	7.66	7.70
Electrical conductivity <sup>a</sup>	28600	19900	8650	5290
Total solids <sup>b</sup>	46300	31090	22380	15890
Total dissolved solids <sup>b</sup>	36250	16930	11565	6420
Total suspended solids <sup>b</sup>	10360	6031	5119	1930
Settleable solids <sup>b</sup>	9690	4260	3390	2840
COD <sup>b</sup>	40820	19190	9998	3010
BOD <sup>b</sup>	15880	6960	4285	2620
Carbonate <sup>b</sup>	Nil	Nil	Nil	Nil
Bicarbonate <sup>b</sup>	12800	7030	3320	1120
Total Phosphorous <sup>b</sup>	39.20	23.39	16.20	9.97
Total Potassium <sup>b</sup>	7200	4590	2990	1860
Calcium <sup>b</sup>	920	602	391	203
Magnesium <sup>b</sup>	1552.68	892.19	201.3	101.6
Sulphur <sup>b</sup>	75.2	35.6	18.9	9.9
Sodium <sup>b</sup>	502	296	218	172
Chlorides <sup>b</sup>	6122	3829	3212	2868
Iron <sup>b</sup>	7.9	6.2	3.4	2.3
Manganese <sup>b</sup>	1020	829	442	201
Zinc <sup>b</sup>	1.5	0.98	0.59	0.51
Copper <sup>b</sup>	0.272	0.201	0.092	0.056
Cadmium <sup>b</sup>	0.005	0.003	0.002	0.001
Lead <sup>b</sup>	0.15	0.09	0.07	0.014
Chromium <sup>b</sup>	0.05	0.021	0.01	0.007
Nikel <sup>b</sup>	0.08	0.049	0.03	0.011
Ammonical Nitrogen <sup>b</sup>	744.7	332.42	274.4	155.09
Carbohydrates <sup>c</sup>	21.64	11.32	7.93	5.92

Units: a-µS, b- mg/L, c-%, PTSW- Primary treated distillery Spentwash

**Table 3:** Amount of N, P, K and S (Nutrients) in distillery Spentwash

Chemical parameters	PTSW	1:1 PTSW	1:2 PTSW	1:3 PTSW
Ammonical Nitrogen <sup>b</sup>	744.7	332.42	274.4	155.09
Total Phosphorous <sup>b</sup>	39.20	23.39	16.20	9.97
Total Potassium <sup>b</sup>	7200	4590	2990	1860
Sulphur <sup>b</sup>	75.2	35.6	18.9	9.9

Unit: b- mg/L, PTSW- Primary treated distillery spentwash

**Table 4:** Proximate principles of Maize at different irrigations (in %)

	Raw water	1:1 PTSW	1:2 PTSW	1:3 PTSW
Crude protein (CP)	4.9	3.6	5.3	8.5
Neutral detergent fibre (NDF)	52.0	57.7	50.3	42.4
Acid detergent fibre (ADF)	31.4	35.6	30.6	27.7
Total digestible nutrient (TDN)	58.8	55.6	59.4	61.6

**Table 5:** Proximate principles of Bajra at different irrigations (in %)

	Raw water	1:1 PTSW	1:2 PTSW	1:3 PTSW
Crude protein (CP)	4.9	3.1	6.8	11.5
Neutral detergent fibre (NDF)	59.1	61.9	55.1	49.4
Acid detergent fibre (ADF)	49.7	48.1	41.3	38.2
Total digestible nutrient (TDN)	45.5	46.3	51.4	53.7

**Table 6:** Proximate principles of Oats at different irrigations (in %)

	Raw water	1:1 PTSW	1:2 PTSW	1:3 PTSW
Crude protein (CP)	5.3	4.1	8.4	12.3
Neutral detergent fibre (NDF)	52.8	58.4	51.1	43.1
Acid detergent fibre (ADF)	36.1	40.2	33.1	29.3
Total digestible nutrient (TDN)	53.7	52.2	57.5	60.4

**3. Results and Discussion**

Characteristics of experimental soils such as pH, electrical conductivity, the amount of organic carbon, available nitrogen(N), phosphorous(p), Potassium(K), sulphur(S), exchangeable calcium(Ca), Magnesium(Mg), Sodium(Na), DTPA Iron(Fe), Manganese(Mn), Copper(Cu) and Zinc(Zn) were analysed and tabulated (Table-1). It was found that the soil composition is fit for the cultivation of plants, because it fulfils all the requirements for the growth of plants. Chemical composition of PTSW, 1:1, 1:2 and 1:3 SW such as pH, Electrical conductivity, Total solids(TS), Total dissolved solids(TDS), Total suspended solids(TSS), Settable solids(SS), Chemical oxygen demand(COD), Biological oxygen demand(BOD), carbonates, bicarbonates, Total phosphorous(P), Total potassium (K), Ammonical Nitrogen(N), Calcium(Ca) Magnesium(Mg), Sulphur(S), Sodium(Na), Chlorides(Cl), Iron(Fe), Manganese (Mn), Zinc(Zn), Copper(Cu), Cadmium(Cd), Lead(Pb), Chromium(Cr) and Nickel(Ni), were analysed and tabulated (Table-2). Amount of N, P, K and S contents are presented in Table-3. The proximate principles for quality forage: Crude protein, Neutral detergent fibre, Acid detergent fibre and Total digestible nutrient of all plants were very good in 1:3 spentwash as compared to 1:1, 1:2 and raw water irrigations. However, nutrients uptakes were high in 1:3 than in all other types of irrigations for both plants and there was no negative impact of spentwash on the quality of forage crops on livestock digestibility (Table 4, 5 and 6).

**4. Conclusion**

It was noticed that the nutrients uptake for the three cereal forage crops was largely influenced in case of 1:3 and 1:2 diluted spent wash irrigation than with raw water and 1:1. But in 1:3 spent wash irrigation it showed more uptake of nutrients when compared to 1:2 and 1:1 diluted spent wash in all the tested forage crops. This concludes that, the treated soil is enriched with the plant nutrients such as nitrogen, potassium and phosphorous. It further concludes that, the subsequent use of diluted spent wash for irrigation enriches the soil fertility and hence the diluted spent wash (1:3) is effective eco-friendly irrigation medium for cultivation of cereal forage crops and also it fulfils all the plant constituents for the production of quality forages without any adverse effect to environment.

**5. Acknowledgement**

The authors are thankful to The Nijaveedu Sugars Ltd., Koppa, Maddur Tq. Karnataka, for providing spentwash and Zonal Agricultural Research Station, University of Agricultural Science, V.C.Farm, Mandya for providing forage seeds, lab for soil, spentwash and proximate principles of forages samples analysis.

**6. References**

1. Amar B, Singh Ashisk Biswas, Sivakoti Ramana. Effect of distillery effluent on plant and Soil enzymatic activities and ground nut quality. *J Plant Nutri. Soil Sci.* 2003; 166:345-347.
2. Basavaraju HC, Chandraju S. Impact of distillery spent wash on the nutrients of Leaves vegetables: An Investigation. *Asian J Chem.* 2008; 20(7):5301-5310.
3. Basavaraju HC, Chandraju S. An Investigation of Impact of distillery spent wash on the nutrients of Top Vegetables. *Int J Agri. Sci.* 2008; 4(2):691-696.
4. Chares S. Vinasse in the fertilization of sugarcane 1985; 1:20.
5. Chandraju S, Basavaraju HC. Impact of distillery spent wash on seed germination and growth of leaves Vegetables: An investigation. *Sugar Journal (SISSTA).* 2007; 38:20-50.
6. Chandraju S, Basavaraju HC, Chidankumar CS. Investigation of impact of Irrigation of distillery spent wash on the growth, yield and nutrients of leafy vegetable. *Chem.Env. Res.* 17. (1&2): in press, 2008.
7. Chandraju S, Basavaraju HC, Chidankumar CS. Investigation of impact of Irrigation of distillery spent wash on the nutrients of cabbage and mint leaf. *Indian Sugar* 2008; 39:19-28.
8. Chandraju S, Basavaraju HC, Chidankumar CS. Investigation of impact of Irrigation of distillery spent wash on the nutrients of pulses. *Asian J Chem.* 2008; 20(8):6342-6348.
9. Chandraju S, Basavaraju HC, Chidankumar CS. Investigation of impact of Irrigation of distillery spentwash on the nutrients of some condiments and root vegetables. *Chem Env Res.* 318 *Sugar Tech* 2008; 10(4):314-318
10. Deverajan L, Oblisami G. Effect of distillery effluent on soil fertility status, yield and quality of rice. *Madras Agri J.* 1995; 82:664- 665.
11. Devarajan L, Rajanan G, Ramanathan G, Oblisami G. Performance of field crops under distillery effluent irrigations, *Kisan world* 1994; 21:48-50.

12. Joshi HC, Kalra N, Chaudhary A, Deb DL. Environmental issues related with distillery effluent utilization in agriculture in India, Asia Pac J Environ. Develop. 1994; 1:92-103.
13. Kaushik K, Nisha R, Jagjeeta K, Kaushik CP. Impact of long and short term irrigation of a sodic soil with distillery effluent in combination with bio-amendments. Bioresource Technology 2005; 96(17):1860-1866.
14. Kuntal Hati M, Ashis Biswas K, Kalikinkar Bandypadhyay, Misra K. Effect of post-methanation effluent on soil physical properties under a soybean-wheat system in a vertisol. J Plant Nutri. Soil Sci. 2004; 167(5):584-590.
15. Narayan TR, Dabadgho PM. Forage crops of India, ICAR, New Delhi, 1972.
16. Whyte RO. Grassland and fodder resources of India, ICAR, New Delhi, 1964.
17. Patil JD, Arabatti SV, Hapse DG. A review of some aspects of distillery spent wash (vinase) utilization in sugar cane, Bartiya sugar May, 1987, 9-15.
18. Pathak H, Joshi HC, Chaudhary A, Chaudhary R, Kalra N, Dwivedi MK. Distillery effluent as soil amendment for wheat and rice. J Indian Soc. Soil Sci. 1998; 46:155-157.
19. Pujar SS. Effect of distillery effluent irrigation on growth, yield and quality of crops. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Dharwad, 1995.
20. Ramadurai R, Gearard EJ. Distillery effluent and downstream products, SISSTA, Sugar Journal. 1994; 20:129-131.
21. Ramana S, Biswas AK, Kundu S, Saha JK, Yadava RBR. Physiological response of soybean (*Glycine max L.*) to foliar application of distillery effluent. Ann. Plant Soil Res 2000; 2:1-6.
22. Singh Y, Raj Bahadur. Effect of application of distillery effluent on maize crop and soil properties. Indian J Agri Sci. 1998; 68:70-74.
23. Rajendran K. Effect of distillery effluent on the seed germination, seedling growth, chlorophyll content and mitosis in *Helianthus Annuus*-Indian Botanical Contactor 1990; 7:139-144.
24. Sahai R, Jabeen S, Saxena PK. Effect of distillery waste on seed germination, seedling growth and pigment content of rice. Indian J Eco. 1983; 10:7-10.
25. Samuel G. The use of alcohol distillery waste as a fertilizer, Proceedings of International American Sugarcane Seminar, 1986, 245- 252.
26. Zalawadia NM, Raman S, Patil RG. Influence of diluted spent wash of sugar industries application on yield and nutrient uptake by sugarcane and changes in soil properties. J Indian Soc Soil Sci. 1997; 45:767-769.
27. Eyini M, Jayakumar M. Pannirselvam S. Distillery effluent induced changes in apiculture wax deposits of *Eichornia crassipes*. Indian Journal of Economy. 1990; 20:1-4.