



## **Chemical compositions and the phytochemical constituents of the seed of *Sesamum indicum* grown at Katsina state, northern Nigeria**

**Datti Y<sup>1</sup>, Bayero AS<sup>2</sup>, Salihu I<sup>3</sup>, Isa BK<sup>4</sup>, Musbahu L<sup>5</sup>, Abubakar SA<sup>6</sup>, Nafisa AM<sup>7</sup>, Nura T<sup>8</sup>, Dikko MA<sup>9</sup>**

<sup>1-7</sup> Department of Chemistry, Yusuf Maitama Sule University, Kano, Nigeria

<sup>8</sup> Department of Chemistry, School of Science Education, FCE (T) Bichi, Nigeria

<sup>9</sup> Department of Science Laboratory Technology, Federal Polytechnic Nasarawa, Nigeria

### **Abstract**

The seeds of *Sesamum indicum* (sesame) are one of the most oldest sauce, relish and spice known to man. This study determines the chemical composition (both proximate and the mineral contents), as well as the phytochemical constituent of the variety of sesame seed available in Katsina State, Nigeria. The proximate and phytochemical analysis were carried out using standard procedures, while the mineral content was determined by atomic absorption spectrophotometry. The results for the proximate content of the analysed sesame seeds are moisture 7.63%; crude protein 21.34%; crude fibre 9.54%; crude lipid 41.84%; ash 9.01% and carbohydrate 10.64%. The seeds were also found to contain some minerals including Na 0.37%; K 1.69%; Mg 0.32%; Ca 1.12%; P 0.76%; Zn 0.0073%; Cu 0.038%; Mn 0.0026 and Fe 0.021%. The results for the phytochemical analysis revealed the presence of alkaloids, carbohydrates, flavonoids, phytosterols, proteins, saponins and terpenoids, while tannins were found to be absent. The present findings suggest that sesame seeds can be a good source of nutrients in the diet and may have health and economic benefits due to its fibre and minerals content, as well as its phytochemical contents.

**Keywords:** *Sesamum indicum*, phytochemicals, proximate analysis, mineral content

### **Introduction**

Man cannot live independently without interacting with plants, for his life and survival would be impossible without plants and plant products (Emmanuel *et al.*, 2015) [13]. The plant kingdom is considered as the treasure house for food, potential drugs, other materials and to some extent shelter. The use of traditional medicines based on the compounds derived and/or isolated from plants is a common activity through out the world, and according to World Health Organisation medicinal plants would be the best source to obtain variety of drugs (Thite *et al.*, 2013) [38]. Recent reports have revealed that quite a large number of indigenous plant species are high in nutrients value (Lalas and Tsaknis 2002). However, despite the identification of many of such plants with potentials of serving as food and drugs, there is very little information about their chemical compositions (Blessing *et al.*, 2010) [8]. One of such plants that has played a major role in human/livestock food and drugs since ancient times is the sesame (Olaleye *et al.*, 2015) [30], and it has been recommended as a potential source to solve the problem of micronutrients deficiencies in modern day nutrition (Blessing *et al.*, 2010) [8].

Widely distributed across India and Africa, *Sesamum indicum* L. (Pedaliaceae), also known as sesame is reputed for its folk medicine (Jefferson 2003). Incidentally, sesame seeds are the oldest sauce, relish and spice known to man, dating back to as early as 1600 BC (Nagendra *et al.*, 2012) [26]. The concoction or infusion from the roots, leaves or seeds of sesame have been recommended by traditional health practitioners in Nigeria for the treatment of various ailments including bruised, catarrh, eye pains, inflamed membranes of the mouth, chicken pox, measles and as a hair

shampoo against *Taenia capitis* (Gill 1992) [18]. Several studies have been conducted on different species of sesame grown in the Western part of Nigeria, however there is scanty scientific information on *S. indicum* grown in the northern part of Nigeria. At the beginning of this study there was very few reports on the phytochemical, proximate and mineral composition of the seeds or other parts (leaves and root) of *S. indicum* grown in the study area. It is in view of that the present study was aimed to provide information on the mineral compositions, proximate composition and phytochemical constituents of the seed of *Sesamum indicum* produced in Katsina, North-Western Nigeria.

### **Materials and Methods**

#### **Sample Collection and Preparation**

Dried sesame seed were obtained from three different farms in Dutsin-Ma Local Government of Katsina State, Nigeria. The seeds were milled with pestle and mortar and then packed in an air tight container and stored in a dessicator (containing silica gel) ready for further analysis. All the chemicals used were of analytical grade.

#### **Determination of Mineral Content**

Mineral analysis was determined using 10 g of the milled sesame seed and subjected to dry ashing for 5 hr in well cleaned porcelain crucibles at 550°C. The residue ash was dissolved in 5 ml of HNO<sub>3</sub>/HCl (1:2) and heated gently on a hot plate until brown fumes disappeared and white coloration was formed. The solution on each crucible was filtered into 100 ml volumetric flask and the volume made up to 100 ml with deionized water.

The cations (Na, K, Mg, Ca, P, Zn, Cu, Mn and Fe) were determined using flame atomic absorption spectrophotometer (Shahid *et al.*, 1999) [33].

#### Determination of Proximate Composition

The proximate compositions of the milled sesame seed were determined using standard analytical methods. All measurements were done in triplicates and values presented in percentage.

#### Moisture Content

Two grams of the milled sesame seed was oven-dried in a crucible at 105 °C overnight. The dried sample was then cooled in desiccator for 1 hr and weighed to a constant weight and the percentage loss in weight was expressed as percentage moisture content (AOAC, 1999).

#### Ash Content

The residue remaining was weighed after the ashing of 2 g dried milled sesame seed in a crucible. The ashing was done in a muffle furnace of temperature 550 °C for 6 hr. The ashed sample was cooled in a desiccator and weighed. The percentage residual weighed was expressed as ash content (AOAC, 1999).

#### Crude Lipid Content

Continuous extraction of lipid was done for 5 hr with petroleum ether in a soxhlet extractor with 2.00 g of the sample used to determine the crude lipid content (Udo and Oguwele, 1986) [39].

#### Crude Protein Content

Kjeldahl method was used to determine total protein. Here 1 g of the sample was put into a filter paper and put into a Kjeldahl flask, 10 cm<sup>3</sup> of concentrated H<sub>2</sub>SO<sub>4</sub> were added and digested in a fume cupboard until the solution becomes colorless. The distillation was carried out with 15 mL of 50% of NaOH. The tip of the condenser was dipped into a conical flask containing 6 cm<sup>3</sup> of 4% boric acid in a mixed indicator until a green coloration was observed. Titration was done in the receiver flask with 0.01 M HCl until the solution turned red (Gabriel *et al.*, 2018) [16].

#### Crude Fibre Content

Estimation of the crude fibre was done by acid and alkaline digestion methods 2.00 g of each sample were used with 20% H<sub>2</sub>SO<sub>4</sub> and NaOH solution (Gabriel *et al.*, 2018) [16].

#### Carbohydrate Content

The carbohydrate content of the test sample was determined by estimation using the arithmetic difference method (De Conto *et al.*, 2011; James, 1995) [4, 23].

%CHO =

100 - (% Moisture + % Fat + % Ash + % Fibre + % Protein)

#### Phytochemical Screening

The qualitative phytochemical screening of the sesame seed extracts was carried out according to different standard procedures to ascertain the phytochemical composition of the seed.

#### Test for Alkaloids

Mayer's Test: Two drops of Mayer's reagent were added to 2 ml of the milled sesame seed extract along the sides of test tube (Evans 1997).

#### Test for Carbohydrates

Benedict's Test: Benedict's reagent (0.5 ml) was added to 0.5 ml of the milled sesame seed filtrate. The mixture was heated on a boiling water bath for 2 minutes (Benedict, 1908).

#### Test for Oils and Fats

Spot Test: A small quantity of extract is pressed between two filter papers (Sahira and Cathrine 2015) [32].

#### Test for Flavonoids

Alkaline Reagent Test: A measure of 2 ml of 2% NaOH was mixed with aqueous milled sesame seed extract, a concentrated yellow color was produced which became colorless when 2 drops of the diluted acid was added (Sofowora 1993) [35].

#### Test for Phytosterols

Libermann-Burchard's Test: The milled sesame seed extract (50 mg) was dissolved in of 2 ml acetic anhydride, and to this 2 drops of concentrated sulphuric acid were slowly added along the sides of the test tube (Finar 1986) [15].

#### Test for Proteins

The extract (10 mg) was dissolved in 5 ml of distilled water and filtered through Whatmann No. 1 filter paper and the filtrate was subjected to Biuret test for proteins. Here 2 ml of the filtrate was treated with 1 drop of 2% copper sulphate solution, and 1 ml of (95%) ethanol was added, followed by excess of potassium hydroxide pellets (Gahan 1984) [17].

#### Test for Saponins

The milled sesame seed extract (50 mg) was diluted with distilled water and made up to 10 ml. The suspension is shaken in a graduated cylinder for 15 minutes (Kokate 1999) [23].

#### Test for Terpenoids

Noller's Test: The milled sesame seed extract (2 mg) was taken in a dry test tube and was treated with a bit of tin foil and 0.5 ml of thionyl chloride, followed by gentle heating (Sourabh *et al.*, 2014) [36].

#### Results

The results for the different proximate composition (the sample of the milled sesame seed) including moisture content, ash content, crude lipid content, crude protein content, crude fibre content and carbohydrate content as determined using different standard techniques are presented in Table 1. Again the sample of the milled sesame seed was tested for the presence of some mineral elements including Na, K, Mg, Ca, P, Zn, Cu, Mn and Fe, and the result is presented in Table 2. Lastly, the sample was also tested for different phytochemical composition including alkaloids, carbohydrates, oils and fats, flavonoids, phytosterols, proteins, saponins and terpenes and the results are presented in Table 3.

**Table 1:** Proximate Composition (%) of *Sesamum indicum* Seed

S.no	Component	Mean Value (%)
1.	Moisture	7.63
2.	Crude Protein	21.34
3.	Crude Fibre	9.54
4.	Crude Lipid	41.84
5.	Ash	9.01
6.	Carbohydrate	10.64

**Table 2:** Mineral Content (%) of the Seed of *Sesamum indicum*

Mineral	Na	K	Mg	Ca	P	Zn	Cu	Mn	Fe
Concentration	0.37	1.69	0.32	1.12	0.76	0.0073	0.038	0.0026	0.021

**Table 3:** Qualitative Analysis of the *Sesamum indicum* Seed Extract

S/NO	Phytochemical Constituents	Watermelon Seed Extract
1.	Alkaloids	+
2.	Carbohydrates	+
3.	Flavonoids	+
4.	Phytosterols	+
5.	Proteins	+
6.	Saponins	+
7.	Terpenoids	+
8.	Tannins	-

+ Present - Absent

## Discussion

Result for the proximate constituent analysis of sesame (*Sesamum indicum*) seeds is presented in Table 1, while that of the minerals composition is presented in Table 2. Reports by different researches reveal the proximate compositions of sesame seeds. The moisture content of the sesame analysed was found to be 7.63% and this was found to be within the range of results (4.11%-10.91%) reported by different researchers (Blessing *et al.*, 2010)<sup>[8]</sup> (10.91%); Nagendra *et al.*, 2012<sup>[26]</sup> (6.61%); Makinde and Akinoso 2013 (5.41%); Haftom *et al.*, 2015<sup>[19]</sup> (4.11%). Therefore the values obtained in this study are found to be within the acceptable limits. The low moisture contents observed in the whole plant may enable this plant to possess a long storage capability. Similarly, the percentage protein in the sesame seed was found to be 21.34% and this is supported by reports from Makinde and Akinoso (2013)<sup>[25]</sup> and Haftom *et al.*, (2015)<sup>[19]</sup> who reported a percentage protein content of 26.79% and 22.58% respectively. This means that sesame seed can serve as a good source of this important nutrient, since protein is an essential component of the diet needed for survival of both humans and animals humans (Pugalenthi *et al.*, 2004).

The sesame seed analyzed was found to contain 9.54% of crude fibre, and this was found to be within the range of 3.2% - 10.0% reported by Obianjunwa *et al.*, (2005)<sup>[19]</sup> and El-Khier *et al.*, (2008)<sup>[12]</sup>. Fiber in the diet is important as it helps to maintain human health by reducing blood cholesterol and glucose level in the body Bello *et al.*, (2008)<sup>[7]</sup>. The presence of an appreciable amount fibre in sesame seed makes it a possible remedy for lowering of cholesterol levels in the blood and reduce risk of various cancers (Gabriel *et al.*, 2018)<sup>[16]</sup>. It is also known to expand the inside walls of the colon, easing the passage of waste, thus making it effective against constipation (Betty *et al.*, 2016)<sup>[7]</sup>.

The crude lipid or oil content of the seed extracted from the sesame seed was found to be 41.84% and this was

found to be a little below (but still within) that reported by Haftom *et al.*, (2015) who reported a range of 47.18%-58.52%, with Tashiro *et al.*, (1990) reporting the oil content in the range of 43.4% to 58.8% for 42 strains of sesame with the highest oil content found in white-seeded strain, while Bahkali *et al.*, (1998) reported oil content in Saudi and Indian sesame seeds ranging from 43.2% to 54.0%. The variation in the oil content may be due to variation in variety, soil type, climatic, maturity of plant, the harvesting time of the seeds and the extraction method used (Egbekun and Ehieze 1997; Rahman *et al.*, 2007). Oil provides concentrated energy in diet and enhanced palatability (Hassan *et al.*, 2008)<sup>[20]</sup>.

The carbohydrate content in the sesame seed analyzed in this study was found to be 10.64% and this supported by reports from other researchers (Blessing *et al.*, (2010)<sup>[8]</sup>; Nagendra *et al.*, (2012)<sup>[26]</sup>; Alege *et al.*, (2013)<sup>[29]</sup>; Makinde and Akinoso (2013)<sup>[16]</sup> and Haftom *et al.*, (2015)<sup>[19]</sup>, who reported a near similar results. However, this value is less than the RDA by FAO/WHO (55%), thus a diet of sesame seeds should be taken along side another food with a much higher carbohydrate in order to have the daily required carbohydrate intake Carbohydrates not only provide the most easily accessible energy source for our body, they also play an important role in many physiological processes.

The ash content of the sample analyzed was found to be 9.01%. This result is similar to the results reported by Blessing *et al.*, (2010)<sup>[8]</sup> who reported 9.62%; Makinde and Akinoso (2013)<sup>[25]</sup> who reported 7.31% and Haftom *et al.*, (2015) who reported 9.00%. Sample with appreciable ash content is expected to have high concentration of various mineral elements, which are expected to speed up metabolic processes, improve growth and development (Betty *et al.*, 2016)<sup>[7]</sup>.

The mineral content of the sesame seed analyzed is presented in Table 2, with potassium (K) having the highest value of 1.69%, followed by calcium (Ca) with 1.12%, then phosphorous (P), sodium (Na) and magnesium (Mg) with values of 0.76%, 0.37% and 0.32% respectively, while iron (Fe), copper (Cu), manganese (Mn) and zinc (Zn) have 0.021%, 0.038%, 0.0026% and 0.0073% respectively. This result is supported by the findings reported by Blessing *et al.*, (2010)<sup>[8]</sup>; Nagendra *et al.*, (2012)<sup>[26]</sup>; Alege *et al.*, (2013); Makinde and Akinoso (2013)<sup>[25]</sup> and Haftom *et al.*, (2015)<sup>[19]</sup>. Copper helps the body form collagen and absorbs iron, and plays a role in energy production. Zinc plays role in wound healing as well as treatment to diarrhea. Iron is a mineral that serves several important functions, its main function being to carry oxygen throughout our bodies and making red blood cells (Beard and Dawson 1997). Due to the reasonable concentrations of calcium, magnesium and zinc in the sesame seed, it should be recommended as an element of bone and teeth strengthening, and also as a potential ingredient in cosmetics for stimulation and revitalization. Potassium and sodium help in regulating the water balance and the acid-base balance in the blood and tissues. Therefore sesame seed can easily be used to supplement some of these minerals in the human body.

The phytochemical analysis/screening of the sesame seeds showed the presence of alkaloids, carbohydrates, flavonoids, phytosterols, proteins saponins and terpenoids, while tannins were absent. This result resemble that reported by

Blessing *et al.*, (2010)<sup>[8]</sup> and Nwobasi and Attamah (2017). However presence of tannins was reported by both the group of reseachers which contradicts the findings of this study. Saponins have been reported to be effective and responsible for the treatment of many conditions such as inflammation, pre-and post-menopausal symptoms (Bombardelli and Gabetta, 2001)<sup>[21]</sup>, cardiovascular and hypertension (Yao *et al.*, 2005)<sup>[27]</sup>. Some alkaloids from plant sources are reported to of medicinal actions as analgesics, antispasmodics, anti-cholinergics and anaesthetics (Okwu, 2004)<sup>[19]</sup>, antimicrobial activity of some alkaloids have also been reported (Olaleye, 2007)<sup>[34]</sup>. Flavonoids are used as natural antioxidants in foods and pharmaceutical drugs due to their ability to scavenge reactive oxygen species (Bombardelli and Gabetta, 2001)<sup>[29]</sup>.

## Reference

1. Alege GO, *et al.* The Morphological, Proximate and Mineral Responses of Sesame to Different Nutrient Sources *Global Journal of Bio-Science and Biotechnology*. 2013; 2(1):12-16
2. AOAC (Association of Official Analytical Chemists). Official Methods of Analysis AOAC International Methods 934.01, 988.05, 920.39 and 942.05. Arlington, VA, USA: AOAC International, 1990.
3. Bahkali AH, *et al.* Protein and Oil Composition of Sesame Seeds (*Sesamum indicum*, L.) Grown in the Gizan Area of Saudi Arabia. *International Journal of Food Sciences and Nutrition*. 1998; 49:409-414.
4. Beard JL, *et al.* Iron, in: O'Dell BL. Sunder RA Editors Handbook of Nutritionally Essential Mineral Elements, New York, CRC Press, 1997, 275-334
5. Bello MOI, *et al.* Studies on the Chemical Composition and Antinutrients of Some Lesser Known Nigerian Fruits. *African Journal of Biotechnology*. 2008; 7:3972-3979.
6. Benedict SR. *A Reagent For the Detection of Reducing Sugars* *J Biol. Chem.* 2002; 5(6):485-487.
7. Betty T, *et al.* Watermelon Seeds as Food: Nutrient Composition, Phytochemicals and Antioxidant Activity *International Journal of Nutrition and Food Sciences*. 2016; 5(2):139-144
8. Blessing M, *et al.* Chemical Composition of *Sesamum indicum* L. (Sesame) Grown in Southe-Eastern Nigeria and the Physicochemical Properties of the Seed Oil *Seed Science and Biotechnology*. 2010; 1(4):1-12.
9. Bombardelli E, Gabetta B. Soya Extract, Process for its Preparation and Pharmaceutical Composition. U.S Patent 6280777, 2001.
10. De Conto LC, *et al.* Characterisation of Crude 62 Watermelon Seed Oil by Two Different Extraction Methods. *Journal of American Oil Chemists' Society*. 2011; 88:1709-1714.
11. Egbekun MK, *et al.* Proximate Composition and Functional Properties of Full Fat and Defatted Beniseed (*Sesamum indicum* L.) Flour. *Plant Foods for Human Nutrition*. 1997; 51:35-41.
12. El Khier, *et al.* Chemical Composition and Oil Characteristics of Sesame Seed Varieties Grown in Sudan. *Research Journal of Agriculture and Biological Sciences*. 2008; 14:1-6.
13. Emmanuel CC, *et al.* Traditional Medicine and the Future of Medicinal Plants in Nigeria *Journal of Medicinal Plants Studies*. 2015; 3(4):23-29
14. Evans WC. *Trease and Evans Pharmacognosy*, Harcourt Brace and Company. Asia PVT. Ltd.Singapore, 1997, 131
15. Finar IL. *Stereo Chemistry and the Chemistry of Natural Products*, Longman. 1996; 2:135.
16. Gabriel AF, *et al.* Characterization of Seed Oil from *Citrullus lanatus* (Watermelon) *Direct Res. J. Public Health and Environ. Technol.* 2018; 3(2):34-40.
17. Gahan PB. *Plant Histochemistry and Cytochemistry: An Introduction*. Academic Press, Florida, U.S.A, 1984, 176.
18. Gills LS. *Ethnomedical Uses of Plants in Nigeria*, UNIBEN Press, Nigeria, 1992, 212.
19. Haftom Z, Geremew B, Solomon A. Physico-Chemical Properties of Sesame (*Sesamum indicum* L.) Varieties Grown in Northern Area, *Ethiopia Agricultural Sciences*. 2005; 6:238-246
20. Hassan LG, *et al.* Proximate, Minerals and Anti-nutritional Factors of Daniellia oliveri seed kernel. *Chemclass Journal*. 2008; 5:31-36.
21. Hassan LG, Umar KJ. Proximate and Mineral Compositions of Seeds and Pulp of *Parkia biglobosa* L. *Nigeria Journal of Basic and Applied Sciences*. 2004; 13:15-27
22. James CS. *Analytical Chemistry of Food*. Champman and Hall, London, 1995, 64-65.
23. Kokate CK. *Practical pharmacognosy* 4th edition, Vallabh Prakashan Publication, New Delhi, India, 1999, 82
24. Lalas S, Tsaknis J. Characterization of Moringa oleifera Seed Oil Variety Periyakulam-1. *Journal of Food Composition and Analysis*. 2002; 15:65-77
25. Makinde FM, Akinoso R. Nutrient composition and effect of processing treatments on anti nutritional factors of Nigerian sesame (*Sesamum indicum* Linn) *International Food Research Journal*. 2013; 20(5):2293-2300.
26. Nagendra PM, *et al.* A Review on Nutritional and Nutraceutical Properties of Sesame *J Nutr Food Sci*. 2012; 2(2):1-6
27. Nwobasi CS, Attamah CG. Proximate Analysis and Phytochemical Properties of Sesame (*Sesamum Indicum* L.) Seeds Grown and Consumed In Abakaliki, Ebonyi State, Nigeria *International Journal of Health and Medicine*. 2017; 2(4):1-4
28. Obiajunwa ET, *et al.* Determination of Essential Minerals and Trace Elements in Nigerian Sesame Seeds Using TXRF Technique. *Pakistan Journal of Nutrition*. 2005; 4:393-395.
29. Olaleye MT. Cytotoxicity and Antibacterial Activity of Methanolic Extract of *Hibiscus sabdariffa*. *J. Med. Plants Res.* 2007; 1(1):9-13.
30. Olaleye OO, *et al.* Extraction, Physicochemical and Phytochemical Characterization of Oil from Sesame Seed *Asian Food Science Journal*. 2015; 1(4):1-12.
31. Rahman MS, *et al.* Studies on the Characterization, Lipids and Glyceride Composition of Sesame (*Sesamum indicum* L.) Seed Oil. *Bangladesh Journal of Science and Indian Research*. 2007; 42:67-74.
32. Sahira KB, *et al.* General Techniques Involved in Phytochemical Analysis *International Journal of*

- Advanced Research in Chemical Science. 2015; 2(4):25-32
33. Shahid I, *et al.* Effect of Root Stock on leaf Mineral Composition and Productivity of Kinnow Mandarin. *Int. J. Agric. Biol.* 1999; 1:91-93.
  34. Singh BB, *et al.* Inheritance of Alectra Resistance in Cowpea Genotype B 301. *Crop Science.* 1993; 33:70-72
  35. Sofowora A. *Phytochemical Screening of Medicinal Plants and Traditional Medicine in Africa*, Spectrum Books Ltd, Ibadan, Nigeria, 1993, 151
  36. Sourabh J, *et al.* Preliminary Phytochemical, Pharmacognostical and Physico Chemical Evaluation of Cedrus deodara Heartwood *Journal of Pharmacognosy and Phytochemistry.* 2014; 3(1):91-95
  37. Tashiro T, *et al.* Oil and Minor Components of Sesame (*Sesamum indicum*) Strains. *Journal of the American Oil Chemists' Society.* 1990; 67:508-511.
  38. Thite SV, Chavan YR, Aparadh VT, Kore BA. Preliminary Phytochemical Screening of Some Medicinal Plants. *IJPCBS;* 2013; 3(1):87-90.
  39. Udo EJ, Oguwele JA. *Laboratory Manual for the Analysis of Soil, Plants and Water Samples.* 3<sup>rd</sup> Edition, Department of Crop Production, University of Ilorin, Kwara State Nigeria, 1986, 131-152.
  40. Weaver LT. Feeding the Weanling in the Developing World: Problems and Solutions. *International Journal of Food Science and Nutrition.* 1994; 45:127-134
  41. Yao X, *et al.* New Use of Saponin Compound for Treating Cardiovascular Disease. CN Patent 1,562,064., 2005.